

Original Article

Anaesthesia for Emergency Surgical Patients with Co-existing Hypertension: A Clinical Challenge

SY Yakubu^{*}, II Lawal, ST Muhammad, ME Idris, AT Bako.

Department of Anaesthesia, Ahmadu Bello University Teaching Hospital, Zaria, Nigeria

Article Metrics

Date submitted: 21/9/2020

Date Accepted: 3/12/2020

Date Published: Jan. 2021



www.jmbsr.com.ng

^{*}Correspondence: Dr. Saidu Yusuf Yakubu, Department of Anaesthesia, Ahmadu Bello University Teaching Hospital, Zaria, Nigeria.

Email: saiduyakubu@yahoo.com

Tel: +2348037873339

ABSTRACT

Background: Hypertension is a medical condition that may co-exist with any surgical disease. Hypertension is defined as a sustained increase in blood pressure (BP) >140/90 mm Hg in a patient not taking antihypertensive drugs. It has been estimated that the overall prevalence of hypertension in Nigeria is 28.9%. Hypertension becomes a challenge to the anaesthetist when the hypertensive patient presents for an emergency operation.

Methodology: This study was a retrospective audit review of anaesthetic charts, case notes and operation records of adult patients that were operated at the accident and emergency (A&E) theatre of a tertiary hospital in the North-West zone of Nigeria over a 6 months period during 2014. Patient's age, sex, type of surgery and anaesthetic technique were collated. Data obtained were expressed as numbers and percentages.

Results: A total of 182 adult patients had emergency surgery. Females were 95(52.2%). General surgery had the highest number (38.5%) of patients followed by Obstetrics with 32.4%. Gynaecological surgery had the lowest number of patients (5.5%). Forty (22%) of patients operated during the period had high blood pressure (>140/90mmHg). Twelve of the hypertensive patients (30%) had urological operations followed by general surgical procedures with 11 patients (27.5%). Four patients (10%) had gynaecological surgery. General anaesthesia (GA) was administered in 80% of the cases while the remaining 8 patients had spinal anaesthesia.

Conclusion: Twenty two percent of study patients were hypertensive and most of them had urological or general surgical procedures. GA was the preferred technique of anaesthesia during the review period.

Keywords: Anaesthesia, Emergency, Hypertension, Surgical patient

Access to the article



website: <http://www.jmbsr.com.ng>

doi: <https://doi.org/10.46912/jmbsr.xx>

Find our articles on Google Scholar, fully indexed

How to cite this article:

SY Yakubu, II Lawal, ST Muhammad, ME Idris, AT Bako. Anaesthesia for Emergency Surgical Patients with Co-existing Hypertension: A Clinical Challenge. *J Med Bas Sci Res* 2020;1(1):9-14.

INTRODUCTION

Hypertension is a medical condition that may co-exist with any surgical disease. The 2017 American College of Cardiology/American Heart Association (ACC/AHA) guidelines defined hypertension by two levels: 1. Elevated blood pressure (BP) with a systolic pressure (SBP) between 120 and 129 mm Hg and diastolic pressure (DBP) less than 80 mm Hg, and 2. Stage 1 hypertension, with an SBP of 130 to 139 mm Hg or a DBP of 80 to 89 mm Hg.¹

Hypertension is also described as a sustained increase in blood pressure (BP) >140/90 mm Hg in a patient not taking antihypertensive drugs.² Adeloye and colleagues estimated that the overall prevalence of hypertension in Nigeria is 28.9%.³ Any hypertensive patient could present with a surgical disease requiring an operation. Hypertension becomes a challenge to the anaesthetist when the hypertensive patient presents for an emergency surgical procedure. Therefore before an emergency operation it is paramount to control elevated blood pressure to avoid excessive bleeding and other complications such as myocardial infarction and stroke.

Uncontrolled hypertension constitutes a high risk to anaesthesia and surgery. Peri-operative hypotension is common in hypertensives and this increases the risk of complications.⁴ A study by Monk et al.⁵ documented that intra-operative hypotension but not hypertension is associated with a higher mortality in hypertensive patients undergoing non-cardiac surgery.

Planning of the surgical procedure to a more suitable and safe time is the option of choice in the uncontrolled hypertensive patient presenting for elective surgery. This allows for time to adequately treat and prepare the patient for the conduct of safe anaesthesia. Such time is not available in an emergency and the anaesthetist is faced with the challenge of coming-up with a suitable technique to anaesthetize the patient. We aim to determine the number of emergency hypertensive surgical patients presenting in our healthcare facility. This will assist us to train and re-train ourselves and the younger ones to look out for high blood pressure during the preoperative assessment of patients and administer the appropriate anaesthetic technique and medications to improve patient outcome.

METHODOLOGY

This study was a retrospective audit review of anaesthetic charts,

case notes and operation records of patients that were operated at the accident and emergency (A&E) theatre of a tertiary hospital in the North-West region of Nigeria over a 6 months period in the year 2014. Patient's age, sex, type of surgery and anaesthetic technique were collated. Data obtained were expressed as numbers and percentages.

RESULTS

A total number of 182 adult patients had emergency surgery during the study period. Females constituted 95 (52.2%) of patients. General surgery had the highest number with 70 (38.5%) of patients followed by obstetric surgery which had 59 (32.4%) patients. Gynaecological surgery had the lowest number of 10 (5.5%) patients as shown in Table 1. Forty (22%) of patients operated during the period had high blood pressure (>140/90mmHg). Twelve of the hypertensive patients (30%) had urological operations followed by general surgical procedures with 11 patients (27.5%). Four patients (10%) had gynaecological surgery (Figure 1). The technique of anaesthesia used for the hypertensive patients are as shown in Figure 2.

Table 1. Distribution of 182 adult emergency surgical patients operated by sex

S/No	Surgical specialty	Male (%)	Female (%)	Total No. of patients (%)
1	General surgery	49 (70.0)	21 (30.0)	70 (38.5)
2	Obstetrics	0 (0.0)	59 (100)	59 (32.4)
3	Orthopaedics	18 (81.8)	4 (18.2)	22 (12.1)
4	Urology	20 (95.2)	1 (4.8)	21 (11.5)
5	Gynaecology	0 (0.0)	10 (100.0)	10 (5.5)
	Total	87 (47.8)	95 (52.2)	182 (100.0)




Figure 1. Distribution of 40 emergency surgical hypertensive patients operated




Figure 2. Techniques of Anaesthesia for emergency surgical hypertensive patients

DISCUSSION

Forty (22%) out of the 182 adult patients that had emergency surgery during the period of the study presented with high blood pressure (BP>140/90mmHg). This is an indication that a good number of our surgical patients for emergency operations do present with hypertension. This is also a reflection of the prevalence rate of hypertension in the general community which is 29%.³ Studies have reported that blood pressure tends to increase with age.⁶

Studies from developed countries have shown that the incidence of preoperative hypertension is between 10%-25%.⁷ Improvement in the life expectancy among Africans coupled with dwindling economic situations will make more hypertensive's to be seen as emergency surgical patients. This signifies more challenge to the anaesthetist as he prepares to manage such emergency hypertensive surgical patient. In a similar study conducted in southern Ethiopia by Mengistu et al they documented a preoperative hypertension rate of 6.9% among elective surgical patients. They stated that this is particularly lower than the rates documented in the literature. They attributed the low rate to the difference in lifestyle in Ethiopia compared to other parts of the world.⁸ The strength of their work lies in being prospective in nature.

Postponement of the surgical procedure to a more suitable and safe time is the option of choice in the uncontrolled elective hypertensive surgical patient. This allows for time to adequately prepare the patient for the safe conduct of anaesthesia. However, such time and luxury is not available in the emergency situation and the anaesthetist is faced with the challenge of coming-up with a suitable technique of anaesthesia for the patient.

Anaesthesia for the hypertensive patient is associated with a lot of risks. Howell et al.⁹ in their work documented that there is a small increase in the incidence of peri-operative major cardiovascular adverse events in hypertensive patients with organ damage or with blood pressure >180/110 mmHg but they cannot say for sure whether postponing surgery to reduce blood pressure overcomes this risk. These highlight the need for the anaesthetist to know how to adequately optimize and manage the hypertensive surgical patient in order to reduce morbidity and mortality.

This audit showed that the emergency surgical patient presenting with high blood pressure could be from any of the surgical sub-specialty. However 57.5% (23 out of 40) of the hypertensive

patients managed in our centre had either urological or general surgical procedures. The anaesthetist needed to be aware of this because his choice of technique could be limited based on the site of the surgery. It is also important to note that >50% (12 out of 21 patients) of all the urology patients managed during the period of study had high blood pressure. This may not be unconnected with the fact that urological patients tend to be older and blood pressure increases with age. The availability of various types of anaesthetic drugs will help the anaesthetist to be more flexible in the choice of anaesthetic technique.

In this study both general and spinal anaesthesia were used for the management of the hypertensive surgical patients that had emergency operations. However, GA was used for most (80%) of the patients. The study by Mengistu and co-workers in Ethiopia documented that GA was used in 62.1 % of operations.⁸ The anaesthetist needs to be meticulous in the choice of medications for GA in the management of the hypertensive surgical patient to avoid upsurge in blood pressure. Hypertensive surges greater than 20% from the baseline are associated with adverse outcomes.¹⁰ Care is needed in the choice of drugs for premedication, induction and maintenance of anaesthesia. There is need to use cardio-stable drugs to ensure cardio protection. Thus glycopyrrolate is preferred to atropine for premedication and propofol is preferred to ketamine for induction of anaesthesia.^{11,12} Anaesthetic manipulations such as endotracheal intubation causes tachycardia and hypertension. This occurs as a reflex response to intubation and increases the risk of anaesthesia especially in the hypertensive patient. These responses could be attenuated using various medications.^{13, 14} Vecuronium and atracurium are more cardio-stable and thus preferred for muscle relaxation than pancuronium bromide which causes tachycardia and hypertension.¹⁵ During the maintenance of anaesthesia, mild increases in blood pressure could be lowered by using inhalational agents like halothane and potent analgesic agents like fentanyl.

The use of RA in the anaesthetic management of the hypertensive surgical patient has the advantage of avoiding the airway. However, uncontrolled hypertensives are more at risk of epidural anaesthesia compared to controlled hypertensive patients and therefore need close monitoring. Marked cardiovascular instability may accompany spinal or epidural

anaesthesia in the form of sudden drop in blood pressure.¹⁶ Our hypertensive patients had good outcome due to careful choice of anaesthetic technique and medications. This study was limited by the fact that it was retrospective. There is need to conduct a prospective study to document the exact prevalence of hypertension among surgical patients in Zaria.

CONCLUSION

Twenty two percent of study patients were hypertensive and most of them had urological or general surgical procedures. GA was the preferred technique of anaesthesia during the period of review.

With precise use of medications, GA or regional anaesthesia (RA) can safely be performed in the hypertensive surgical patient undergoing an emergency operation.

Conflicts of interest

No conflicts of interest to report.

REFERENCES

1. Whelton PK, Carey RM, Aronow WS, Casey DE, Collins KJ, Dennison HC, *et al.* ACC /AHA /AAPA /ABC /ACPM /AGS /APhA /ASH /ASPC /NMA / PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association task force on clinical practice guidelines. *Circulation*. 2018; 138: 484–594.
2. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL *et al.* The Seventh report of the Joint National Committee on Prevention, Detection, Evaluation and Treatment of High Blood Pressure. *Hypertension*. 2003; 42(6):1206-52.
3. Adeloye D, Basquill C, Aderemi AV, Thompson JY, Obi FA. An estimate of the prevalence of hypertension in Nigeria: a systematic review and meta-analysis. *J Hypertens*. 2015; 33:230–42.
4. Walsh M, Devereaux PJ, Garg AX, Kurz A, Turan A, Rodseth RN *et al.* Relationship between intraoperative mean arterial pressure and clinical outcomes after noncardiac surgery: toward an empirical definition of hypotension. *Anesthesiology* 2013; 119: 507–15.
5. Monk TG, Bronsert MR, Henderson WG, Mangione MP, Sum-Ping ST, Bentt DR, *et al.* Association between intraoperative hypotension and hypertension and 30-day postoperative mortality in noncardiac surgery. *Anesthesiology* 2015; 123:307-19.
6. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL *et al.* National heart, lung, and blood institute joint national committee on prevention, detection, evaluation, and treatment of high blood pressure; national high blood pressure education program coordinating committee. The seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure: the JNC 7 report. *JAMA*. 2003; 289:2560–72.
7. Varon J, Marik PE. Perioperative hypertension management. *Vasc Health Risk Manag* 2008; 4:615-27.
8. Mengistu K, Geleta D. Prevalence of preoperative hypertension and its intraoperative anesthetic management among patients undergoing elective surgery at Hawassa University. *Int. J. Med. Med. Sci*. 2019; 11(2):11-19.
9. Howell SJ, Sear JW, Foëx P. Hypertension, hypertensive heart disease and perioperative cardiac risk. *Br J Anaesth* 2004; 92:570-83.
10. Hartle A, McCormack T, Carlisle J, Anderson S, Pichel A, Beckett N, *et al.* The measurement of adult blood pressure and management of hypertension before elective surgery: Joint Guidelines from the Association of Anaesthetists of Great Britain and Ireland and the British Hypertension Society. *Anaesthesia* 2016; 71:326-37.
11. Desalu I, Kushimo OT, Bode CO. Comparative study of the haemodynamic effects of atropine and glycopyrrolate at induction of anaesthesia in children. *West Afr J Med*. 2005; 24(2):115-119.
12. Best W, Bodenschatz C, Beran D. World Health Organisation Critical Review of Ketamine. 36th WHO Expert Committee on Drug Dependence report, 6.2. Geneva, Switzerland: World Health Organisation. 2014.
13. Hassani V, Movassaghi G, Goodarzi V, Safari S. Comparison of fentanyl and fentanyl plus lidocaine on attenuation of hemodynamic responses to tracheal intubation in controlled hypertensive patients undergoing general anesthesia. *Anesth*

Pain Med. 2013; (3):115-118.

14. Lee JH, Kim H, Kim HT, Kim M, Cho K, Lim SH et al. Comparison of dexmedetomidine and remifentanyl for attenuation of hemodynamic responses to laryngoscopy and tracheal intubation. *Korean J Anesthesiol.* 2012; 63(2):124-129.
15. Satwik T, Basavaraj P, Hulkund SY, Ajay BC. Comparison of rocuronium, vecuronium and atracurium in tiva for hemodynamic effects during beating heart bypass surgery. *Indian J Clin Anaesth.* 2019; 6(3); 366-370.
16. Kristensen SD, Knuuti J, Saraste A, Anker S, Bøtker HE, De Hert S, *et al.* 2014 ESC/ESA Guidelines on non-cardiac surgery: Cardiovascular assessment and management. The Joint Task Force on non-cardiac surgery: Cardiovascular assessment and management of the European Society of Cardiology (ESC) and the European Society of Anaesthesiology (ESA). *Eur J Anaesthesiol* 2014; 31:517-73.